PRENATAL NUTRITION MODULE POST-TEST ANSWER KEY

100 Total Points

PRENATAL NUTRITION MODULE POST-TEST REVISED 12/99

		NAME
		DATE
		SCORE
		HE FOLLOWING STATEMENTS T FOR TRUE OR F FOR FALSE. T the false statements.
	1.	THE THIRD TRIMESTER IS THE MOST CRITICAL PHASE OF DEVELOPMENT. IT IS DURING THIS TIME THAT EXPOSURE TO DRUGS AND ALCOHOL IS MOST LIKELY TO LEAD TO BIRTH DEFECTS.
	2.	A WOMAN'S PREGRAVID WEIGHT IS HER WEIGHT JUST BEFORE SHE BECAME PREGNANT.
	3.	WIC STAFF WOULD NEED TO RISK A WOMAN WITH HIGH MATERNAL WEIGHT GAIN IF SHE GAINED 11 POUNDS SINCE HER LAST WIC VISIT 6 WEEKS AGO.
	4.	THE NUMBER OF SERVINGS OF DAIRY PRODUCTS RECOMMENDED FOR THE PREGNANT TEEN IS GREATER THAN THAT FOR THE ADULT PREGNANT WOMAN.
	5.	LISTERIA IS A BACTERIA FOUND IN HARD CHEESES.
	6.	A WOMAN WHO DECIDES TO TAKE AN ADDITIONAL SUPPLEMENT OF VITAMIN C DURING PREGNANCY BECAUSE HER FRIEND TOLD HER IT WOULD HELP KEEP HER HEALTHY SHOULD BE RISK WITH NUTRITION RISK FACTOR 97 "EXCESSIVE INTAKE OF DIETARY SUPPLEMENTS, VITAMINS, OR MINERALS."
	7.	WIC STAFF SHOULD RECOMMEND PRENATAL WIC PARTICIPANTS CUT BACK ON THEIR SALT INTAKE IF THEY REPORT SWELLING DURING THEIR PREGNANCY.
	8.	IT IS PROBABLY SAFE FOR PREGNANT WOMEN TO CONSUME UP TO 300 MILLIGRAMS OF CAFFEINE A DAY.
	9.	ONE DRINK OF ALCOHOL EVERY TWO WEEKS IS PROBABLY SAFE DURING PREGNANCY.
1	10.	WOMEN WHO SMOKE DURING PREGNANCY ARE MORE LIKELY TO GIVE BIRTH TO LOW BIRTH WEIGHT BABIES.

CHECK ALL THAT APPLY

11.	WHICH OF THE FOLLOWING ARE NUTRITION RISK FACTORS FOR				
	THE PREGNANT WIC PARTICIPANT?				
	PICA				
	ANEMIA				
	HIGH PARITY (HISTORY OF 3 OR MORE PREGNANCIES)				
	ELEVATED BLOOD LEAD VALUE				
	OVERWEIGHT 26.0 BMI				
	UNDERWEIGHT 19.8 BMI				
	SHORT STATURE				
	SMOKING				
	MATERNAL AGE _ 35 YEARS				
	MATERNAL AGE _ 17 YEARS				
	CLOSELY SPACED PREGNANCIES				
	MIGRANCY				
	HYPEREMESIS GRAVIDARUM				
	BEGINNING PRENATAL CARE DURING THE 11TH WEEK OF				
	PREGNANCY				
12.	WHAT THREE COMPONENTS OF PREGNANCY ACCOUNT FOR THE				
12.	MAJORITY OF THE WEIGHT GAINED IN PREGNANCY?				
	Mayora i or the werdin drained in the drainer.				
	BABY WEIGHT				
	UTERUS ENLARGEMENT				
	BREAST ENLARGEMENT				
	PLACENTA				
	MOTHER'S FAT STORES				
	BLOOD AND FLUID INCREASES				
13.	WHICH OF THE FOLLOWING ARE NUTRITION RISK FACTORS FOR				
	THE POSTPARTUM, NON-BREASTFEEDING WIC PARTICIPANT?				
	PICA SMOKING				
	ANEMIA MATERNAL AGE _ 35 YEARS				
	HIGH PARITY MATERNAL AGE 17 YEARS				
	HIGH PARITY MATERNAL AGE _ 17 YEARS (HISTORY OF 3 OR CLOSELY SPACE PREGNANCIES				
	MORE PREGNANCIES) MIGRANCY				
	ELEVATED BLOOD LEAD HYPEREMESIS GRAVIDARUM				
	VALUE BEGINNING PRENATAL				
	OVERWEIGHT 26.0 BMI CARE DURING THE 11 TH WEEK				
	UNDERWEIGHT 19.8 BMI OF PREGNANCY				
	SHORT STATURE				

14.	COMPARE THE TWO FOODS, AND SE NUTRIENT DENSE:	LECT THE (ONE THAT IS MORE
	PEANUT BUTTER ON CELERY	OR	CREAM CHEESE ON CELERY
	SODA POP	OR	_ORANGE JUICE
	GLAZED DONUT	OR	_BAGEL WITH CHEESE
	BEAN BURRITO WITH SALSA	OR	BLT (BACON, Lettuce, tomato Sandwich)
	POTATO CHIPS & SOUR CREAM DIP	OR	PRETZELS AND YOGURT
CON	APLETE THE FOLLOWING QUESTIONS		
15.	LIST <u>two</u> reasons why adequate pregnancy is important. 1.	NUTRITIO	ON DURING
	2.		
16.	LIST ONE REASON WHY ANEMIA DUI	RING PREG	NANCY IS A RISK.
17.	DEFINE THE WEEKS FOR EACH OF THE FIRST TRIMESTER SECOND TRIMESTER THIRD TRIMESTER	HE THREE T —	TRIMESTERS:
18.	LIST ONE REASON WHY INADEQUAT FETUS.	E WEIGHT	GAIN IS A RISK TO THI
19.	LIST ONE REASON WHY EXCESSIVE W FETUS.	/EIGHT GA	IN IS A RISK TO THE
20.	LIST ONE REASON WHY EXCESSIVE W PREGNANT WOMAN.	/EIGHT GA	IN IS A RISK TO THE
21.	IDENTIFY THE WEIGHT CATEGORY F	FOR THE FC	OLLOWING FOUR

PREGNANT WOMEN.

					PRY: PRY:		
	JEAN STEVENS B.						
	BERNIE TAOS B						
22.	WHAT ARE THE REE				HT GAIN	ranges fo	R
	MARIA WONG TO SUSAN DIAZ TO JEAN STEVENS TO BERNIE TAOS TOT	OTAL WE	IGHT GAI	IN RANG IN RANG	E: E:		
23.	LIST FOUR QUESTI Underweight Pi Education and 1.	REGNANT	Γ WOMAN	TO HEL			ON
	2.						
	3.						
	4.						
24.	WHAT NUTRIENT DEFECTS?	IS IMPOR	RTANT TO	PREVEN	T NEURAI	. TUBE	
25.	LIST <u>Two</u> nutrie. The vegan diet. 1.	nts tha	T ARE MC	RE LIKEI	LY TO BE I	NADEQUAT	'E IN
	2.						
26.	LIST FOUR TIPS TH Nausea. 1.	HAT MAY	HELP A W	oman d	EAL WITH	i prenatal	,
	2.						
	3.						

	4.
27.	WHAT ARE SIX HEALTHY TIPS WIC STAFF CAN SELECT FROM TO EDUCATE POSTPARTUM WOMEN ON? 1.
	2.
	3.
	4.
	5.
	6.
FIL	L IN THE BLANKS
28.	WHAT ARE THE NUMBER OF SERVINGS FOR THE DIET OF A PREGNANT WOMAN RECOMMENDED BY THE FOOD GUIDE PYRAMID FOR EACH OF THE FOLLOWING FOOD GROUPS:
	DAIRY PRODUCTS GROUP MEAT/MEAT ALTERNATIVES GROUP FRUIT GROUP VEGETABLE GROUP BREADS/CEREALS GROUP
29.	WHAT ARE THE NUMBER OF SERVINGS FOR THE DIET OF A POSTPARTUM NON-BREASTFEEDING WOMAN RECOMMENDED BY TH FOOD GUIDE PYRAMID FOR EACH OF THE FOLLOWING FOOD GROUPS:
	DAIRY PRODUCTS GROUP MEAT/MEAT ALTERNATIVES GROUP FRUIT GROUP VEGETABLE GROUP BREADS/CEREALS GROUP

PRENATAL NUTRITION MODULE POST-TEST ANSWER KEY

100 POINTS TOTAL

MARK THE FOLLOWING STATEMENTS T FOR TRUE OR F FOR FALSE. CORRECT THE FALSE STATEMENTS. (2PTS EACH)

F	1. THE THIRD [FIRST] TRIMESTER IS THE MOST CRITICAL PHASE OF DEVELOPMENT. IT IS DURING THIS TIME THAT EXPOSURE TO DRUGS AND ALCOHOL IS MOST LIKELY TO LEAD TO BIRTH DEFECTS.
<u>T</u>	2. A WOMAN'S PREGRAVID WEIGHT IS HER WEIGHT JUST BEFORE SHE BECAME PREGNANT.
<u>T</u>	3. WIC STAFF WOULD NEED TO RISK A WOMAN WITH HIGH MATERNAL WEIGHT GAIN IF SHE GAINED 11 POUNDS SINCE HER LAST WIC VISIT 6 WEEKS AGO.
<u>T</u>	4. THE NUMBER OF SERVINGS OF DAIRY PRODUCTS RECOMMENDED FOR THE PREGNANT TEEN IS GREATER THAN THAT FOR THE ADULT PREGNANT WOMAN.
F	5. LISTERIA IS A BACTERIA FOUND IN HARD [SOFT] CHEESES.
<u>T</u>	6. A WOMAN WHO DECIDES TO TAKE AN ADDITIONAL SUPPLEMENT OF VITAMIN C DURING PREGNANCY BECAUSE HER FRIEND TOLD HER IT WOULD HELP KEEP HER HEALTHY SHOULD BE RISK WITH NUTRITION RISK FACTOR 97 "EXCESSIVE INTAKE OF DIETARY SUPPLEMENTS, VITAMINS, OR MINERALS."
<u> </u>	7. WIC STAFF SHOULD [NOT] RECOMMEND PRENATAL WIC PARTICIPANTS CUT BACK ON THEIR SALT INTAKE IF THEY REPORT SWELLING DURING THEIR PREGNANCY. [THEY SHOULD REFER PARTICIPANTS TO THEIR HEALTH CARE PROVIDER.]
<u>T</u>	8. IT IS PROBABLY SAFE FOR PREGNANT WOMEN TO CONSUME UP TO 300 MILLIGRAMS OF CAFFEINE A DAY.
<u>F</u> _	9. ONE DRINK OF ALCOHOL EVERY TWO WEEKS IS PROBABLY SAFE DURING PREGNANCY. [THERE IS NO KNOWN SAFE AMOUNT OF ALCOHOL FOR A HEALTHY PREGNANCY.]
<u>T</u>	10. WOMEN WHO SMOKE DURING PREGNANCY ARE MORE LIKELY TO GIVE BIRTH TO LOW BIRTH WEIGHT BABIES.

CHECK ALL THAT APPLY

11.	WHICH OF THE FOLLOWING ARE NUTRITION RISK FACTORS FOR THE
	PREGNANT WIC PARTICIPANT? (14PTS)
T	_ PICA
T	_ ANEMIA
	_ HIGH PARITY (HISTORY OF 3
	OR MORE
	PREGNANCIES)
T	_ ELEVATED BLOOD LEAD
	VALUE
	_ OVERWEIGHT 26.0 BMI
<u></u>	_ UNDERWEIGHT 19.8 BMI
	_ SHORT STATURE
<u>T</u>	_ SMOKING
	_ MATERNAL AGE _ 35 YEARS
	_ MATERNAL AGE _ 17 YEARS
<u></u>	
	PREGNANCIES
<u></u>	_ MIGRANCY
<u></u>	_ HYPEREMESIS GRAVIDARUM
	_ BEGINNING PRENATAL CARE
	DURING THE
	11TH WEEK OF PREGNANCY
12.	WHAT THREE COMPONENTS OF PREGNANCY ACCOUNT FOR THE MAJORITY OF THE WEIGHT GAINED IN PREGNANCY? (3PTS)
13.	WHICH OF THE FOLLOWING ARE NUTRITION RISK FACTORS FOR THE
	POSTPARTUM, NON-BREASTFEEDING WIC PARTICIPANT? (14PTS)
<u></u>	PICA SHORT STATURE
<u></u>	ANEMIA SMOKING
	HIGH PARITY (HISTORY OF 3 MATERNAL AGE _ 35 YEARS
	OR MORE PREGNANCIES) <u>T</u> MATERNAL AGE _ 17 YEARS
<u></u>	
	VALUE PREGNANCIES
<u></u>	OVERWEIGHT 26.0 BMI <u>T</u> MIGRANCY
T_	UNDERWEIGHT 19.8 BMI HYPEREMESIS GRAVIDARUM

	BEGINNING PRENATAL CARE DURING THE 11 th	WEEK OF PREGNANCY
14.	COMPARE THE TWO FOODS, AND SELECT NUTRIENT DENSE: (5PTS)	T THE ONE THAT IS MORE
	T PEANUT BUTTER ON CELERY	OR CREAM CHEES ON CELERY
	SODA POP	OR <u>T</u> ORANGE JUICE
	GLAZED DONUT	OR <u>T</u> BAGEL WITH CHEESE
		OR BLT (BACON, LETTUCE, TOMATO SANDWICH)
	POTATO CHIPS & SOUR CREAM DII	P OR <u>T</u> PRETZELS AND YOGURT
CON	APLETE THE FOLLOWING QUESTIONS	
15. (2PT		ELOPMENT OF THE FETUS.
	2. FOR A HEALTHY PREGNANCY A	ND PREGNANCY OUTCOME.
16. (1PT	LIST ONE REASON WHY ANEMIA DURIN THE FETUS WOULD NOT RECEIVE ADEC GROWTH. ANEMIA DURING PREGNAN DELIVERY OF LOW BIRTH WEIGHT INF RISK OF INFANT MORTALITY.	QUATE OXYGEN FOR PROPER ICY IS ASSOCIATED WITH THI
17. (3PT	DEFINE THE WEEKS FOR EACH OF THE	THREE TRIMESTERS:
	FIRST TRIMESTER SECOND TRIMESTER THIRD TRIMESTER 27-40	
18.	LIST ONE REASON WHY INADEQUATE V The Fetus.	WEIGHT GAIN IS A RISK TO
(1PT) THE INFANT MAY BE BORN WITH A LO	W BIRTH WEIGHT AND WITH

LIST ONE REASON WHY EXCESSIVE WEIGHT GAIN IS A RISK TO THE

FETAL GROWTH RESTRICTION.

19.

FETUS.

- (1PT) THE INFANT MAY BE BORN WITH HIGH BIRTH WEIGHT. HIGH BIRTH WEIGHT INFANTS RISK INJURY AT DELIVERY.
- 20. LIST ONE REASON WHY EXCESSIVE WEIGHT GAIN IS A RISK TO THE PREGNANT WOMAN.
- (1PT) EXCESSIVE WEIGHT GAIN IS ASSOCIATED WITH OTHER COMPLICATIONS OF PREGNANCY INCLUDING HIGH BLOOD PRESSURE, PREECLAMPSIA, AND ECLAMPSIA.
- 21. IDENTIFY THE WEIGHT CATEGORY FOR THE FOLLOWING FOUR PREGNANT WOMEN.

(4PTS)

MARIA WONG	BMI 20.7	WEIGHT CATEGORY: NORMAL
SUSAN DIAZ	BMI 28.2	WEIGHT CATEGORY: <u>High</u>
JEAN STEVENS	BMI 19.0	WEIGHT CATEGORY: <u>Low</u>
BERNIE TAOS	BMI 32.0	WEIGHT CATEGORY: <u>Obese</u>

22. WHAT ARE THE RECOMMENDED TOTAL WEIGHT GAIN RANGES FOR EACH OF THE WOMEN IN QUESTION 21?

(4PTS)	MARIA WONG	TOTAL WEIGHT GAIN RANGE : <u>25-35 Pounds</u>
	SUSAN DIAZ	TOTAL WEIGHT GAIN RANGE : 15-25 POUNDS
	JEAN STEVENS	TOTAL WEIGHT GAIN RANGE : 28-40 POUNDS
	BERNIE TAOS	TOTAL WEIGHT GAIN RANGE : <u>15 Pounds</u>

23. LIST FOUR QUESTIONS YOU COULD ASK OF AN OVERWEIGHT OR UNDERWEIGHT PREGNANT WOMAN TO HELP IDENTIFY NUTRITION EDUCATION AND COUNSELING NEEDS?

(4PTS) EXAMPLES ARE LISTED BELOW. OTHERS ARE ACCEPTABLE AS APPROPRIATE.

HOW DO YOU FEEL ABOUT GAINING WEIGHT?
HOW IS YOUR APPETITE?
WHAT TYPE OF PHYSICAL ACTIVITIES DO YOU ENGAGE IN?
DO YOU HAVE ENOUGH FOOD RESOURCES?
HOW MUCH WEIGHT ARE YOU THINKING YOU SHOULD GAIN FOR THIS PREGNANCY?

24. WHAT NUTRIENT IS IMPORTANT TO PREVENT NEURAL TUBE DEFECTS?

(1PT) FOLATE

- 25. LIST <u>Two</u> nutrients that are more likely to be inadequate in the vegan diet.
- (2PTS) ANY TWO OF THESE: VITAMIN B₁₂, CALCIUM, IRON, VITAMIN D

26. LIST FOUR TIPS THAT MAY HELP A WOMAN DEAL WITH PRENATAL NAUSEA. ANY FOUR OF THE FOLLOWING, OR ADDITIONAL ONES, IF (4PTS) APPROPRIATE: SMELL/TASTE LEMON SIP LIQUIDS FREQUENTLY FRESH AIR DRY CEREAL, TOAST, **CRACKERS** AVOID STRONG-SMELLING FOODS SEVERAL SMALL MEALS. **RATHER THAN 3 LARGE MEALS EAT SLOWLY** PREPARE COLD FOODS BE RELAXED WHEN EATING AVOID FOODS THAT **UPSET YOUR STOMACH** WHAT ARE SIX HEALTHY TIPS WIC STAFF CAN SELECT FROM TO 27. EDUCATE POSTPARTUM WOMEN ON? 1. EAT RIGHT (6PTS) 2. EAT FOODS RICH IN FOLATE EVERY DAY 3. **BE ACTIVE** SEE A HEALTH CARE PROVIDER 4. MAKE TIME FOR BEING A NEW MOM 5. STAY SMOKE-FREE 6. FILL IN THE BLANKS 28. WHAT ARE THE NUMBER OF SERVINGS FOR THE DIET OF A PREGNANT WOMAN RECOMMENDED BY THE FOOD GUIDE PYRAMID FOR EACH OF THE FOLLOWING FOOD GROUPS: <u>4</u> (5 FOR TEEN) DAIRY PRODUCTS GROUP (5PTS)MEAT/MEAT ALTERNATIVES GROUP 3 FRUIT GROUP 2-4 VEGETABLE GROUP 3-5 BREADS/CEREALS GROUP 6-11 29. WHAT ARE THE NUMBER OF SERVINGS FOR THE DIET OF A POSTPARTUM NON-BREASTFEEDING WOMAN RECOMMENDED BY THE FOOD GUIDE PYRAMID FOR EACH OF THE FOLLOWING FOOD **GROUPS:** DAIRY PRODUCTS GROUP <u>3</u> (4 FOR TEEN) (5PTS)MEAT/MEAT ALTERNATIVES GROUP FRUIT GROUP

VEGETABLE GROUP